arXiv:1408.7032v1 [math.CO] 25 Aug 2014

The Bounds for Eigenvalues of Normalized and Signless Laplacian Matrices

ŞERİFE BÜYÜKKÖSE AND ŞEHRİ GÜLÇİÇEK ESKİ

June,2014

In this paper, we obtain the bounds of the extreme eigenvalues of a normalized and signless Laplacian matrices using by their traces. In addition, we determine the bounds for k-th eigenvalues of normalized and signless Laplacian matrices.

AMS classification: 05C50, 05C75

Keywords: Normalized Laplacian matrix, Signless Laplacian matrix, Bounds of eigenvalue.

1 Introduction

Let $G(V,E)$ be a simple graph with the vertex set $V=\{v_1,v_2,\ldots,v_n\}$ and edge set of E. For $v_i \in V$, the degree of v_i, the set of neighbours of v_i denoted by d_i and N_i, respectively. The cardinality of N_i is denoted by c_{ij}; i.e, $|N_i| = c_{ij}$. If v_i and v_j adjacency, we denote $v_i \sim v_j$ of shortly use $i \sim j$.

The adjacency matrix, Laplacian matrix and diagonal matrix of vertex degree of a G graph denoted by $A(G)$, $L(G)$, $D(G)$, respectively. Clearly

$$L(G) = D(G) - A(G).$$

The normalized Laplacian matrix of G is defined as $\mathscr{L}(G) = D^{-1/2}(G)L(G)D^{-1/2}(G)$ i.e, $\mathscr{L}(G) = [\ell_{ij}]_{n\times n}$, where

$$\ell_{ij} = \begin{cases} 1 & ; \quad if \quad i=j \\ \frac{-1}{\sqrt{d_i d_j}} & ; \quad if \quad i \sim j \\ 0 & ; \quad otherwise. \end{cases}$$

The signless Laplacian matrix of G is defined as $Q(G) = D(G) + A(G)$ i.e, $Q(G) = [q_{ij}]_{n\times n}$, where

$$q_{ij} = \begin{cases} d_i & ; \quad if \quad i=j \\ 1 & ; \quad if \quad i \sim j \\ 0 & ; \quad otherwise. \end{cases}$$

Since $\mathscr{L}(G)$ normalized Laplacian matrix and $Q(G)$ signless Laplacian matrix are real symetric matrices, their eigenvalues are real. We denote the eigenvalues of $\mathscr{L}(G)$ and $Q(G)$ are by

$$\lambda_1(\mathscr{L}(G)) \geq \cdots \geq \lambda_n(\mathscr{L}(G))$$

and

$$\lambda_1(Q(G)) \geq \cdots \geq \lambda_n(Q(G))$$

,respectively.

Now we give some bounds for normalized Laplacian matrix and signless Laplacian matrix.

Oliveira and de Lima's bound [1]. For a simple connected graph G with n vertices and m edges, $\Delta = d_1 \geq d_2 \geq \cdots \geq d_n = \delta$

$$\lambda_1(Q(G)) \leq \max_i \left\{ \frac{d_i + \sqrt{d_i^2 + 8d_i m_i}}{2} \right\} \tag{1}$$

where $m_i = \frac{1}{d_i} \sum_{i \sim j} d_j$.

Another Oliveira and de Lima's bound [1].

$$\lambda_1(Q(G)) \leq \max_i \left\{ d_i + \sqrt{d_i m_i} \right\} \tag{2}$$

where $m_i = \frac{1}{d_i} \sum_{i \sim j} d_j$.

Li, Liu et al. bound's [2,3].

$$\lambda_1(Q(G)) \leq \frac{\Delta + \delta - 1\sqrt{(\Delta + \delta - 1)^2 + 8(2m - (n-1)\delta)}}{2}. \tag{3}$$

Rojo and Soto's bound [4]. If λ_1 is the largest eigenvalue of $\mathscr{L}$, then

$$|\lambda_1(\mathscr{L}(G))| \leq 2 - \min_{i<j} \left(\frac{|N_i \cap N_j|}{\max\{d_i, d_j\}} \right) \tag{4}$$

where the minimum is taken over all pairs (i, j), $(1 \leq i < j \leq n)$.

In this paper, we find an extreme eigenvalues of normalized Laplacian matrix and signless Laplacian matrix of a G graph with using their traces.

To obtain bounds for eigenvalues of $\mathscr{L}(G)$ and $Q(G)$ we need the following lemmas and theorems.

Lemma 1. Let W and $\lambda = (\lambda_j)$ be nonzero column vectors, $e = (1, 1, \ldots, 1)^T$, $C = I_n - \frac{ee^T}{n}$ $m = \frac{\lambda^T e}{n}$, $s^2 = \frac{\lambda^T C \lambda}{n}$ and I_n is an Identity matrix. Let $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$. Then

$$-s\sqrt{nW^TCW} \leq W^T\lambda - mW^Te = W^TC\lambda \leq s\sqrt{nW^TCW}.$$

$$\sum_j (\lambda_j - \lambda_n)^2 = n[s^2 + (m - \lambda_n)^2]$$

$$\sum_j (\lambda_1 - \lambda_j)^2 = n[s^2 + (\lambda_1 - m)^2].$$

$$\lambda_n \leq m - \frac{s}{\sqrt{n-1}} \leq m + \frac{s}{\sqrt{n-1}} \leq \lambda_1.$$

Theorem 1. Let A be a $n \times n$ complex matrix. Conjugate transpose of A denoted by A^*. Let $B = AA^*$ whose eigenvalues are $\lambda_1(B) \geq \lambda_2(B) \geq \cdots \geq \lambda_n(B)$. Then

$$m - s\sqrt{n-1} \leq \lambda_n^2(B) \leq m - \frac{s}{\sqrt{n-1}}$$

and

$$m + \frac{s}{\sqrt{n-1}} \leq \lambda_1^2(B) \leq m + s\sqrt{n-1}$$

where $m = \frac{trB}{n}$ and $s^2 = \frac{trB^2}{n} - m$.

2 Main Results for Normalized Laplacian Matrix

Theorem 2. Let G be a simple graph and $\mathscr{L}(G)$ be a normalized Laplacian matrix of G. If the eigenvalues of $\mathscr{L}(G)$ are $\lambda_1(\mathscr{L}(G)) \geq \lambda_2(\mathscr{L}(G)) \geq \cdots \geq \lambda_n(\mathscr{L}(G))$, then

$$\lambda_n(\mathscr{L}(G)) \leq \sqrt{\left(1 + \frac{2}{n} \sum_{i\sim j, i<j} \frac{1}{d_i d_j}\right) + \sqrt{\frac{tr[L(G)]^4 - nm^2}{n(n-1)}}} \tag{5}$$

$$\lambda_1(\mathscr{L}(G)) \geq \sqrt{\left(1 + \frac{2}{n} \sum_{i\sim j, i<j} \frac{1}{d_i d_j}\right) + \sqrt{\frac{tr[L(G)]^4 - nm^2}{n(n-1)}}} \tag{6}$$

$$\lambda_1(\mathscr{L}(G)) \leq \sqrt{1 + \frac{2}{n} \sum_{i\sim j, i<j} \frac{1}{d_i d_j} + \sqrt{\left(\frac{tr[L(G)]^4}{n} - m^2\right)(n-1)}} \tag{7}$$

Proof. Obviously,

$$tr[\mathscr{L}(G)]^2 = n + 2 \sum_{i\sim j, i<j} \frac{1}{d_i d_j}$$

and

$$tr[\mathscr{L}(G)]^4 = \sum_{i=1}^{n} \left(1 + \sum_{i\sim j} \frac{1}{d_i d_j}\right)^2 + 2 \sum_{i<j} \left(\sum_{k \in N_i \cap N_j} \frac{1}{d_k \sqrt{d_i d_j}} - \sum_{i\sim j} \frac{2}{\sqrt{d_i d_j}}\right)^2$$

Since $\mathscr{L}(G)$ real symmetric matrix, we find the result from Theorem 1.

Example 1. Let $G = (V, E)$ with $V = \{1, 2, 3, 4, 5, 6\}$ and $E = \{(1,2), (1,5), (2,3), (2,4), (2,6), (3,4), (3,5), (4,5), (5,6)\}$.

$\lambda(\mathscr{L}(G))$	(4)	(6)(lower bound)	(7)(upper bound)
1.86	2	1.34	1.93

3 Main Results for Signless Laplacian Matrix

Theorem 3. Let G be a simple graph and $Q(G)$ be a signless Laplacian matrix of G. If the eigenvalues of $Q(G)$ are $\lambda_1(Q(G)) \geq \lambda_2(Q(G)) \geq \cdots \geq \lambda_n(Q(G))$, then

$$\lambda_n(Q(G)) \leq \sqrt{\left(1 + \frac{2}{n} \sum_{i \sim j, i<j} \frac{1}{d_i d_j}\right) + \sqrt{\frac{tr[Q(G)]^4 - nm^2}{n(n-1)}}} \tag{8}$$

$$\lambda_1(Q(G)) \geq \sqrt{\left(1 + \frac{2}{n} \sum_{i \sim j, i<j} \frac{1}{d_i d_j}\right) + \sqrt{\frac{tr[Q(G)]^4 - nm^2}{n(n-1)}}} \tag{9}$$

$$\lambda_1(Q(G)) \leq \sqrt{1 + \frac{2}{n} \sum_{i \sim j, i<j} \frac{1}{d_i d_j} + \sqrt{\left(\frac{tr[Q(G)]^4}{n} - m^2\right)(n-1)}} \tag{10}$$

Proof. Clearly

$$tr[Q(G)]^2 = n + 2 \sum_{i \sim j, i<j} \frac{1}{d_i d_j}$$

and

$$tr[Q(G)]^4 = \sum_{i=1}^{n} \left(1 + \sum_{i \sim j} \frac{1}{d_i d_j}\right)^2 + 2 \sum_{i<j} \left(\sum_{k \in N_i \cap N_j} \frac{1}{d_k \sqrt{d_i d_j}} - \sum_{i \sim j} \frac{2}{\sqrt{d_i d_j}}\right)^2$$

Since $Q(G)$ real symmetric matrix, we found the result from Theorem 1.

Example 2. Let $G = (V, E)$ with $V = \{1, 2, 3, 4, 5, 6, 7\}$ and
$E = \{(1,2), (1,3), (1,4), (1,5), (1,6), (1,7), (2,3), (3,5), (4,5), (4,6)\}$.

$\lambda(Q(G))$	(1)	(2)	(3)	(9)(lower bound)	(10)(upper bound)
7.67	9.08	9.74	9.34	4.58	7.76

REFERENCES

[1] C.S. Oliveira, L.S.De Lima, N.M.M. De Abreu, P. Hansen, *Bound on the Index of the Signless Laplacian of a graph, Discrete Appl. Math.* **158**, (2010), 355-360

[2] J. Li, Y. Pan, *Upper Bounds for the Laplacian Graph Eigenvalues, Acta Math. Sin. Engl. ser. 20(5)*, (2004), 803-806

[3] H. Liu, M. Lu, F. Tian, *On the Laplacian Spectral Radius of a Graph, Linear Algebra Appl.*, (2004) ,**376**, 135-141

[4] O. Rojo, R.L. Soto, *A New Upper Bound On The Largest Normalized Laplacian Eigenvals, Operators and Matrices*, (2013) number2, volume7, 323-332

[5] H. Wolkowich, G.P.H. Styan, *Bounds for Eigenvalues Using Traces of Matrice*, *Linear Algebra and its Applications*, (1980), **29**, 471-506

Şerife BÜYÜKKÖSE
Gazi University Faculty of Science
Department of Mathematics
06500 Ankara-Turkey
e-mail: sbuyukkose@gazi.edu.tr
serifebuyukkose@gmail.com

Şehri Gülçiçek ESKİ
Ahi Evran University
Institue of Science
40100 Kırşehir-Turkey
e-mail: gulcicekeski@gmail.com